

University of Dundee

Effectiveness of nutritional and exercise interventions to improve body composition and muscle strength or function in sarcopenic obese older adults

Theodorakopoulos, Christos; Jones, Jacklyn; Bannerman, Elaine; Greig, Carolyn A.

Published in:
Nutrition Research

DOI:
[10.1016/j.nutres.2017.05.002](https://doi.org/10.1016/j.nutres.2017.05.002)

Publication date:
2017

Licence:
CC BY-NC-ND

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Theodorakopoulos, C., Jones, J., Bannerman, E., & Greig, C. A. (2017). Effectiveness of nutritional and exercise interventions to improve body composition and muscle strength or function in sarcopenic obese older adults: A systematic review. *Nutrition Research*, 43, 3-15. <https://doi.org/10.1016/j.nutres.2017.05.002>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Effectiveness of nutritional and exercise interventions to improve body composition and muscle strength or function in sarcopenic obese older adults: A systematic review

Christos Theodorakopoulos¹, Jacklyn Jones^{1*}, Elaine Bannerman², Carolyn A. Greig³

¹Dietetics Nutrition & Biological Sciences, School of Health Sciences, Queen Margaret

University, Edinburgh, United Kingdom, EH21 6UU;

email: CTheodorakopoulos@qmu.ac.uk (C Theodorakopoulos); JJones@qmu.ac.uk (J Jones)

²Centre for Public Health Nutrition Research, School of Medicine, University of Dundee,

Dundee, United Kingdom, DD1 4HN; email: E.Bannerman@dundee.ac.uk

³School of Sport, Exercise and Rehabilitation Sciences, and MRC-Arthritis Research UK

Centre for Musculoskeletal Ageing Research, University of Birmingham, Birmingham,

United Kingdom, B15 2TT; email: C.A.Greig@bham.ac.uk

* Corresponding author

Department of Dietetics, Nutrition and Biological Sciences, School of Health Sciences,

Queen Margaret University, Queen Margaret University Drive, Edinburgh, United Kingdom,

EH21 6UU

18 **List of Abbreviations**

19

20 % BF; Percentage Body Fat

21 ACSM; American College of Sports Medicine

22 AHA; American Heart Association

23 BIA; Bioelectrical Impedance Analysis

24 BMI; Body Mass Index

25 CT; Computerized Tomography

26 DXA; Dual-energy X-ray Absorptiometry

27 EPIDOS; EPIDemiologie de l'OSteoporose

28 EWGSOP; European Working Group on Sarcopenia in Older People

29 FM; fat mass;

30 GFR; Glomerular Filtration Rate

31 HD; Habitual Diet

32 HG, Handgrip

33 HSC; High Speed Circuit

34 IGF-1; Insulin-like Growth Factor 1

35 IADL; Instrumental Activities of Daily Living

36 LM; Lean Mass

37 MRI; Magnetic Resonance Imaging

38 MPS; muscle protein synthesis

39 mTOR; mechanistic Target of Rapamycin

40 PRISMA; Preferred Reporting for Systematic Reviews and Meta-analyses

41 PROSPERO; International prospective register of systematic reviews

- 42 RCH+HD; Ricotta Cheese plus Habitual Diet
- 43 RCTs; Randomized controlled trials
- 44 RPE; Rates of Perceived Exertion
- 45 RM; Repetition Maximum
- 46 SD; Standard Deviation
- 47 SGOT; Serum Glutamic Oxaloacetic Transaminase
- 48 SGPT; Serum Glutamic-Pyruvic Transaminase
- 49 SPPB; Short Physical Performance Battery test
- 50 SMI; Skeletal muscle index; SO, Sarcopenic Obesity
- 51 SH; Strength Hypertrophy
- 52 TASM; Total appendicular skeletal muscle

Abstract

Although sarcopenic obesity (SO) poses a major public health concern, a robust approach for the optimization of body composition and strength/function in SO has not yet been established. The purpose of this systematic review was to assess the effectiveness of nutritional (focusing on energy and protein modulation) and exercise interventions, either individually or combined, on body composition and strength/function in older adults with SO. MEDLINE, the Cochrane Central Register of Controlled Trials, CINAHL and SPORTDiscus were searched. Main inclusion criteria comprised sarcopenia as defined by the European Working Group on Sarcopenia in Older People (EWGSOP) and obesity defined as % body fat ≥ 40 % (women) and ≥ 28 % (men). Randomized controlled trials (RCTs), randomized controlled crossover trials and controlled clinical trials with older adults (mean age ≥ 65 years) following a nutritional regimen and/or an exercise training program were considered. Out of 109 full text articles identified, only two RCTs (61 participants) met the inclusion criteria. One study was a nutritional intervention adding 15 g protein·day⁻¹ (via cheese consumption) to the participants' habitual diet. The second study was a high-speed circuit resistance training intervention. Body composition did not change significantly in either of the studies. However, the exercise intervention improved significantly muscle strength and physical function. Although this review was limited by the small number of eligible studies, it provides evidence for the potential benefits of exercise and highlights the necessity for future research to develop effective interventions including dietary and exercise regimens to combat sarcopenic obesity.

Keywords: Aged; sarcopenia; obesity; dietary proteins; exercise; systematic review

1. Introduction

Sarcopenia is defined by the European Working Group on Sarcopenia in Older People (EWGSOP) as the age-related decline of muscle mass and strength or function [1]. Low strength and muscle mass are associated with poor functional status, physical impairments, frailty, increased risk of falls, loss of independence and higher mortality risk [1][2]. It has been suggested that in older people, strength is a stronger predictor of functional impairment and mortality rates than absolute changes in muscle mass or lean mass alone [3-6]. Secondary to functional impairments, muscle atrophy may also contribute to insulin resistance as muscle tissue plays the main role in glucose uptake and utilization [7]. According to a recent systematic review, the prevalence of sarcopenia may vary from 1 % to 29 % in community-dwellers and 14-33 % in long-term care populations [8].

Another condition that can promote poor health is obesity, which is defined as ‘abnormal or excess body fat accumulation’ [9], and is a growing concern due to its progressively rising prevalence rates in older populations [10]. In 2010, 35 % and 28 % of the adults 65 years of age and older were reported to be obese in the US and the UK, respectively [11][12]. Similar to sarcopenia, obesity can increase the risk of falls and mobility limitations in older age [13][14], and when used in conjunction with indices of body composition and fat distribution (waist circumference or waist to hip ratio) it may be associated with adverse health effects, such as cardiovascular disease, metabolic syndrome, diabetes mellitus and several cancers [15]. Furthermore, adipose tissue can infiltrate the muscle tissue [16] and mediate an inflammatory response [17], which can result in muscle atrophy, mobility losses and lower strength and muscle quality [16][18][19].

100

101 The relationship between sarcopenia and obesity is complex, with the development/
102 progression of one condition being closely connected to the other (**Figure 1**). The condition
103 where sarcopenia and obesity occur together has been termed sarcopenic obesity (SO) [20]. It
104 has been suggested that SO can predispose older individuals to more physical disabilities, gait
105 and balance abnormalities, and an increased risk of falls compared with either of the two
106 conditions alone [21]. Individuals with SO are exposed to ~2.5 times higher risk of reporting
107 Instrumental Activities of Daily Living (IADL) disabilities compared with adults without
108 obesity but with sarcopenia, or adults with obesity but without sarcopenia [22]. This negative
109 synergistic effect of sarcopenia and obesity is in accordance with the findings from the
110 EPIDOS (EPIDemiologie de l'OSteoporose) study, which reported that among a cohort of
111 1,308 women divided in four groups: 1) without sarcopenia or obesity 2) with obesity but not
112 sarcopenia 3) with sarcopenia but not obesity and 4) with SO, the latter was the poorest in
113 terms of performing physical activities that required strength [23]. According to a meta-
114 analysis of 12 prospective cohort studies with a total number of 35,287 participants, the
115 adults with SO had a 24 % higher risk of all-cause mortality compared with their healthy
116 counterparts [24].

117

118 **Figure 1** Relationship between sarcopenia and obesity and associated outcomes

119

120 Although SO has gained significant attention from the scientific community in recent years,
121 and a plethora of existing definitions and cut-offs for sarcopenia and obesity exist, there is no
122 universally accepted definition for SO [1][25][26]. Depending on the definition criteria and

cut-offs used the prevalence rates of SO can vary up to 26-fold, which makes detection and management of the condition challenging for healthcare practitioners [27]. Moreover, there are operational challenges around the management of SO. While exercise training can be beneficial for both obesity and sarcopenia, the dietary management of obesity may require energy restriction, whilst management of sarcopenia requires an increased intake of macronutrients, especially protein [28].

This has resulted in a growing body of evidence highlighting potentially beneficial nutritional and exercise strategies, aiming to reverse or attenuate the negative effects of ageing on body composition and physical function [29-31]. Particular focus has been placed on protein intake, energy modulation and resistance exercise [32][33]. With regard to protein intake, there seems to be a consensus for the benefits of increased protein intake, ranging from 1.0 g·kg bw⁻¹·day⁻¹ to 1.5 g·kg bw⁻¹·day⁻¹, with the higher values appropriate for those older adults with chronic conditions, sarcopenia and malnutrition, or when combined with resistance exercise [28][34][35].

However, there are relatively few intervention studies utilizing exercise training and/or nutritional regimens for older adults with SO [26][36]. It appears that most intervention trials have aimed to attenuate muscle loss at an early stage rather than try to ‘reverse’ an established condition related to advanced ageing such as sarcopenia or SO, which would be far more challenging [37]. Furthermore, to the best of our knowledge, there has been no systematic review to date assessing the effectiveness of nutritional and exercise strategies, alone or combined, to improve body composition and strength/function indices in older individuals with SO. Therefore, the purpose of this systematic review was to assess the

evidence for the use of diets modulating energy and protein (or amino acids) content, exercise training regimens, or diet and exercise training combined, in older adults with SO.

The focus of this systematic review was to determine the effectiveness of protein or energy-modulating regimens, with or without exercise training on body composition and function in adults, 65 years of age and older with SO. In particular, our aims were to 1) determine changes in absolute muscle mass, total appendicular skeletal muscle (TASM), skeletal muscle index (SMI), fat mass, % body fat, body weight and body mass index (BMI), 2) assess changes in muscle strength and/or physical function (including muscle strength, power, gait speed and balance and 3) evaluate the effect of these interventions on quality of life, metabolic profile, activities of daily living, adverse effects of supplementation or food choices, compliance rates and changes in habitual dietary intake during or after the interventions.

2. Approach

This systematic review was performed according to the Preferred Reporting for Systematic Reviews and Meta-analyses (PRISMA) guidelines [38]. The protocol was registered with the International prospective register of systematic reviews (PROSPERO registration number: CRD42015017311).

2.1. Search Strategy

The Cochrane Central Register of Controlled Trials, MEDLINE (via EBSCOhost Research Databases), CINAHL and SPORTDiscus were searched up to and including May 2016. The last search was conducted on 22 May 2016. No limits were applied for date of publication. Combinations of key terms with Medical Subject Headings (MeSH) and Boolean operators were used. The main keywords and terms used were: Age*/ Adult*/ Old*/ Elderly/ Senior, Sarcopeni*/ Lean/ Frail/ Atrophy/ Weakness, Obes*/ Overweight/ Body Mass Index, Exercise/ Training/ Strength/ Muscle/ Mass/ Hypertrophy/ Size/ Body Composition, Diet/ Supplements/ Protein/ Amino Acids/ Energy, Life Quality/ Intervention. The search limiters were English language and studies with human participants [the complete search strategy is presented in supplementary file 2].

2.2. Inclusion Criteria

We included randomized control trials (RCTs), randomized control crossover trials and controlled clinical trials using prospective nutritional and/or exercise interventions to attenuate/ reverse the loss of muscle mass, reduce adipose tissue and optimize muscle strength or function. Given that there are no universally adopted definition criteria for SO, some authors may have used different terms to define the participants, e.g. ‘weak and overweight’ or ‘obese frail’ etc. Such studies were included only if the participants had a sarcopenic phenotype based on the definition criteria and cut-off scores recommended by the EWGSOP [1]. Therefore, studies were included only if they presented data for a) body composition (data on absolute muscle mass, appendicular muscle mass, Total Appendicular Skeletal Muscle (TASM) or Skeletal Muscle Index (SMI) assessed by Dual-energy X-ray Absorptiometry (DXA), Bioelectrical Impedance Analysis (BIA), Computerised Tomography (CT) or Magnetic Resonance Imaging (MRI) and b) muscular strength and/ or physical

function identified by one of the following tests: handgrip strength, knee flexion/extension, peak expiratory flow, gait speed, the Short Physical Performance Battery test (SPPB), the timed up-and-go test or the stair climb power test . The mean age cut-off for inclusion was ≥ 65 years based on how ‘old age’ is defined in the joint recommendations from the American College of Sports Medicine (ACSM) and the American Heart Association (AHA) [39]. The inclusion criterion for obesity was defined as mean percentage body fat (% BF) ≥ 28 % in men and ≥ 40 % in women [22] or in the absence of % BF data, a BMI $\geq 27 \text{ kg}\cdot\text{m}^{-2}$. For any given BMI, a person with sarcopenia will have by definition more body fat compared with their counterparts without sarcopenia, therefore, adults with sarcopenia can present high-adiposity at BMIs substantially lower than $30 \text{ kg}\cdot\text{m}^{-2}$ [40]. Moreover, it is not uncommon for studies to recruit participants who would be classified as overweight or obese based on a BMI cut-off ranging from 25 to $28 \text{ kg}\cdot\text{m}^{-2}$ when the focus is on sarcopenic obesity and/or when participants come from a non-Caucasian ethnic group [25][26]. Studies that presented neither the % BF nor BMI were included only if these indices could be derived from the weight, height and body fat mass values, or if the authors of the study when contacted provided the essential information.

Nutritional interventions aiming to promote muscle hypertrophy by macronutrient profile modification or weight loss via energy restriction were of primary interest. Studies providing extra macronutrients (especially proteins or amino acids and their metabolites) either in the form of whole foods or dietary supplements administered through the oral route only were considered. Exercise regimens including resistance, balance, aerobic and mixed exercise protocols influencing lean mass, fat mass, muscle hypertrophy, strength, power, speed and/ or physical functional were also of primary interest.

219

220 **2.3. Exclusion Criteria**

221

222 Studies were excluded if the protocol involved administration of any kind of prescription
223 only/pharmaceutical agents, or any type of supplementation administered via a route other
224 than oral. Studies including participants with cachexia or with serious mental and cognitive
225 conditions prohibiting adherence to a structured exercise/ nutrition regimen, such as
226 Alzheimer's or dementia, were excluded.

227

228 **2.4. Study Selection**

229

230 The titles and abstracts were screened for eligibility (CT), and the full text copies of
231 potentially eligible articles were obtained for further inspection. The full-text articles were
232 independently assessed for eligibility by two reviewers (CT and JJ). The reference lists of
233 eligible articles and review papers as well as journals specializing in older age were hand
234 searched for potential articles. Any disagreement between the two reviewers was resolved by
235 a third reviewer (CAG).

236

237 **2.5. Data Extraction**

238 Data were extracted from each eligible article by two reviewers (CT and CAG). Any
239 disagreements between the two reviewers were resolved by discussion until consensus was
240 reached. Demographic (age, sex, ethnicity/host country and habitation), methodological
241 (study design, sample sizes, duration, nutritional/dietary and/or exercise intervention plan,
242 supplement type, dosing/frequency of administration, exercise training

type/frequency/volume, assessment method, blinding) and outcome data (changes within and between groups, significance, dropout rates, compliance, adverse effects) were compiled in a standardized Excel spreadsheet.

2.6. Quality Assessment

The quality of the studies was assessed by two independent reviewers using a modified version of the Downs and Black rating scale [41][42]. The Downs and Black scale is one of the most credible instruments for the quality assessment of randomized [43] and non-randomized intervention trials [44]. Modified scoring for Question 27 was performed as detailed by Eng et al. [42]: the original scale had a maximum score of 32 but in this review Question 27 was modified to score either 0 or 1 point instead of the original 0-5 points. Therefore, the maximum total score for the five sections of the scale (reporting, external validity, internal validity/bias, internal validity/confounding, power) was 28.

2.7. Principal Summary Measures

The primary outcome measures were 1) differences in mean of skeletal muscle mass (either absolute, relative or appendicular) and body fat or BMI, and 2) differences in mean of muscle strength and physical function/performance

3. Results

3.1. Description of studies

Our search strategy resulted in 1,440 potential articles. After the exclusion of 1,331 articles based on titles and abstracts, 109 full-text articles reporting 109 studies were retrieved and assessed for eligibility. The detailed flow chart of the selection process is presented in **Figure 2**. The authors of two potentially eligible studies [45][46] were contacted for further information, but retrieval of all the essential body composition data was not possible for reasons unrelated to this review, therefore, the articles were excluded. A total of $n=2$ studies [47][48] including $n=61$ participants met the inclusion criteria and were included in the review. Study A[47] was a nutritional intervention and study B[48] an exercise training intervention; neither of the studies combined exercise with diet.

Figure 2. Information flow through the phases of the systematic review according to PRISMA guidelines.

3.2. Quality Assessment

The two studies were randomized control trials of moderate methodological quality based on the modified Downs and Black rating scale [41][42]. The total score for each study was 18 out of 28. The summary key information of the methodological strengths and limitations is presented in table 1 (**supplementary file 1** presents the complete breakdown of the scoring in the different subsections of the scale). Both studies performed power calculations to determine the population sample size prior to recruitment, however, study B[48] was underpowered; target was $n=21$ per group, but the final analysis was conducted with $n=9$ and $n=8$ for the control and intervention group, respectively. In study A[47] only the testers were

blinded but not the participants. In study B[48] the two groups were exercising at different times, therefore, participants were partially blinded. Study A[47] reported and tested for a range of potential confounders, but failed to report essential information regarding the participants' dietary intake at baseline and follow-up. The results in Study A were based on an intention-to-treat analysis, whereas in Study B the analysis conducted was per-protocol.

Table 1. Summary key points of the included study designs.

3.3. Participant Characteristics

Participants in study A[47] were physically-independent individuals living in Mexico. Their mean \pm SD age and TASM were 76 \pm 5.4 years and 15.5 \pm 2.9 kg, respectively. The mean % BF of men and women was 33.3 \pm 6.2 % and 47.8 \pm 6.6 %, respectively. At baseline two men had a % BF < 28 % and three women < 40 %. All participants in study B[48] were independent-living community dwellers from South Miami (USA). The mean \pm SD age and BMI of participants was 71.3 \pm 7.8 years and 32.6 \pm 4.7 kg \cdot m⁻², respectively and their mean SMI 6.6 \pm 1.0 kg \cdot m⁻².

3.4. Study Design

The aim of study A[47] was to assess whether the addition of a protein rich food to the habitual diet could increase TASM and strength in older individuals with sarcopenia. The study was a 3-month RCT with a control (habitual diet; HD) and an intervention group (habitual diet + 210 g ricotta cheese per day; RCH+HD). The cheese provided 15.7 g extra protein (including 8.6 g of essential amino acids), 10.4 g carbohydrate, 18.4 g fat and a total

of 267 kcal per day. Cheese was divided into three 70 g portions and participants were instructed to consume each portion along with their usual breakfast, lunch and dinner. Dual-energy x-ray absorptiometry was used to measure TASM and body composition changes.

Study B[48] was a 15-week single blind RCT, which aimed to assess the effectiveness of a novel exercise regime based on a high speed circuit (HSC) resistance training program (intervention) on body composition, muscular performance and IADL compared with a conventional strength hypertrophy (SH) regime (control group) in community-dwellers with SO. Body composition was assessed by single frequency BIA. Both groups performed exercises at 11 pneumatic gym machines twice per week. The SH protocol involved three sets of 10-12 repetitions at 70 % of 1RM with a 1-2 min recovery break between sets. Participants were instructed to keep a similar speed of contraction for both the concentric and eccentric phase (2 seconds per phase). The HSC group performed 10-12 repetitions at the same 11 exercises, but in a circuit pattern (i.e. moving from one exercise to the other) with no break in between exercises, unless one full circuit was complete. Three full circuits were performed in total. The resistance load was selected based on maximum power output for each machine. The concentric phase was performed as fast as possible while the eccentric in 2 seconds. No dietary or nutritional element was introduced in the study, and neither dietary patterns nor intakes were reported.

3.5. Outcomes

3.5.1. Body composition

No significant changes were seen in body composition, in either experimental or control groups. In study A[47] the addition of ricotta cheese resulted in no significant changes in lean mass, TASM or body fat in the intervention or control group (Table 2). Secondary analysis by sex showed that although men ($n=8$) in the intervention group experienced an increase in TASM by 490 g, this was not significantly different either from baseline or when compared against the control group ($p=0.42$), which gained a non-significant 220 g of TASM. Similarly, in study B[42] no statistically significant differences were detected in any of the body composition indices, regardless of the exercise regimen (Table 2). Skeletal muscle index (SMI) increased non-significantly in both groups (from $6.5 \pm 0.66 \text{ kg} \cdot \text{m}^{-2}$ to $6.6 \pm 0.59 \text{ kg} \cdot \text{m}^{-2}$ in HSC and from $6.7 \pm 0.45 \text{ kg} \cdot \text{m}^{-2}$ to $6.8 \pm 0.42 \text{ kg} \cdot \text{m}^{-2}$ in SH).

Table 2. Summary of the included studies

3.5.2 Strength and/ or function

In study A[47], the group receiving the extra protein noted a non-significant trend towards an increase in strength (+ 0.9 % relative increase). Although the control group experienced a drop in strength (-3.5 %), the difference between the two groups did not achieve statistical significance ($p=0.06$).

Study B[48] reported significant improvements in several aspects of strength and function in both exercise groups (Table 2). In particular, the strength-hypertrophy (SH) control group experienced significant improvements in leg press 1RM by 22 % ($p<0.01$), chest press 1RM by 16 % ($p=0.03$), leg press peak power by 19 % ($p=0.03$) and chest press peak power by 15% ($p<0.01$) whereas a non-significant increase of 12% was detected in hand grip strength (from $17.3 \pm 2.7 \text{ kg}$ to $19.4 \pm 4.6 \text{ kg}$; $p>0.05$). The HSC group had a significant improvement

in chest press 1RM by 21 % ($p<0.01$), leg press peak power by 41% ($p<0.01$) chest press peak power by 24 % ($p<0.01$) but hand grip strength did not change significantly (increased by 10 %, from 17.7 ± 7.8 kg to 19.4 ± 6.6 kg; $p>0.5$). Between group differences were detected only for leg press peak power, with the HSC group performing better than the control by 158 W [95 % CI (2, 315), $p=0.005$].

The Short Physical Performance Battery (SPPB) test improved significantly over time only within the HSC group from 8.0 ± 1.5 to 9.6 ± 1.2 ($p=0.02$). Between group differences favored the HSC group [mean difference 1.1 (95 % CI (-0.1, 2.4), $p=0.08$], although this was not statistically significant.

3.5.3 Secondary Outcomes

Consumption of ricotta cheese in study A[47], resulted in significantly lower fasting insulin levels in men ($p=0.05$) but not in women. There were no other significant changes in hepatic markers (Serum Glutamic Oxaloacetic Transaminase (SGOT), Serum Glutamic-Pyruvic Transaminase (SGPT) and Alkaline Phosphatase), kidney function (blood urea, uric acid, creatinine and Glomerular Filtration Rate (GFR)), anabolism (Insulin-like Growth Factor-1 (IGF-1)) or insulin resistance. No cases of microalbuminuria were present in the RCH+HD group after the intervention period. In the intervention group, 25 % of women reported early satiety after the consumption of ricotta cheese, however, dietary intakes were not reported. Eight participants from the intervention group dropped out; five were due to personal health issues, two could not eat the entire portion of ricotta cheese, and one had to relocate. In the control group three people dropped out (two for personal reasons and one for modifying the

habitual diet). However, all participants were measured pre- and post-intervention according to an intention-to-treat analysis.

The exercise intervention in study B[48] resulted in acute joint pain only in the SH group. In addition, the HSC group reported significantly lower rates of perceived exertion (RPE) with a mean difference of -1.5 (95 % CI -2.0,-0.12, $p=0.04$). Adherence rates were similar in the two groups; 81 % in HSC and 85 % in SH. Regarding the Instrumental Activities of Daily Living (IADLs) there were significant improvements within both groups (pre vs post); namely, time needed for jacket on and off (from 11.5 ± 3.5 s to 10.2 ± 2.0 s; $p=0.04$), scarf pick-up (5.2 ± 1.1 s to 4.7 ± 0.91 s; $p<0.01$ and pan carry (4.9 ± 0.61 s to 3.9 ± 0.77 s; $p<0.01$) improved significantly within the control group, while the HSC group experienced significant improvements in time for sit-to-stand (from 16.1 ± 5.7 s to 13.4 ± 3.9 s; $p=0.02$) and pan carry (5.4 ± 1.3 s to 4.5 ± 1.2 s; $p<0.01$). No differences were observed between the two groups in the aforementioned IADLs.

In summary, neither of the studies had a significant effect on body composition. The introduction of ricotta cheese in the habitual diet of participants in study A[47] aimed to increase their protein intake but it was not reported whether or not this was achieved, and if so to what extent. In the same study, there was a trend for an increase in strength in the intervention group, but this was not significant. The only significant improvement reported was the fasting insulin levels, but that was reported only in men in the intervention group. Despite the lack of body composition changes, in study B[48], the high speed circuit resistance training and strength hypertrophy resistance training protocols significantly improved strength, power and function indices. Finally, due to the limited data extracted and

diversity of methodologies, statistical pooling was not feasible and therefore, a narrative analysis was conducted.

4. Discussion

The aim of this systematic review was to assess the effectiveness of nutritional and/or exercise interventions on body composition and strength or function in older adults with obesity and sarcopenia. Although only two studies were identified, the lack of intervention trials clearly highlights the need for more research in this area, especially trials combining exercise with nutritional approaches targeting this population group. With regard to the main outcomes, neither an increase in protein intake by $15\text{g} \cdot \text{day}^{-1}$ nor a 15-week resistance exercise protocol produced significant improvements in body composition indices. However, the exercise intervention (both the control group following a strength-hypertrophy resistance exercise protocol and the intervention group utilizing a high-speed power-orientated circuit resistance training) reported significant improvements in both strength and function.

4.1 Effects of protein intake on body composition and function in sarcopenic obesity

Study A[47] attempted to utilize the effects of protein on increased skeletal muscle mass accretion rates. Although the authors acknowledged that the suggested recommendations for protein intake in older individuals with sarcopenia are $1.2\text{--}1.5\text{ g} \cdot \text{kg} \text{ bw}^{-1} \cdot \text{day}^{-1}$ [47], they did not report the participants' daily protein intake, therefore, it was not corroborated whether such intakes were achieved. It has been suggested that maximal muscle protein synthesis (MPS) rates in older adults can be achieved using $\sim 35\text{--}40\text{ g protein} \cdot \text{meal}^{-1}$ [49-51] or $0.4\text{ g protein} \cdot \text{kg} \text{ bw}^{-1} \cdot \text{meal}^{-1}$ [51]. A valid question would be whether a daily addition of 210 g ricotta cheese (delivering 15.7 g protein [47]) to the habitual diet could practically augment

muscle mass in older adults with sarcopenia. It is important to note that the cheese servings were not consumed in one meal, instead they were spread over the three main meals, that is, 70 g cheese (~5 g of extra protein per meal) consumed with breakfast, lunch and dinner. Protein intakes in Study A[47] were not reported, but if we extrapolate data from studies in similar population cohorts [52], it has been suggested that older individuals are not likely to consume an adequate amount of protein during all main meals. Tieland et al. [52] reported mean protein intakes of ~8 g, ~18 g and ~29 g for breakfast, lunch and dinner, respectively. Therefore, it is uncertain whether the addition of 5 g protein in the main meals in study A[47] was enough to significantly augment MPS.

Another confounder may have been the potential impact of the addition of cheese on the habitual diet given the fact that 25 % of women in Study A[47] experienced early satiety. It could be consequently speculated that women's habitual diet was modified with the addition of ricotta, potentially displacing the intake of other foods. However this cannot be confirmed as the habitual diet was not reported.

In study A [47], even though there was a trend towards increased strength, it could be argued that higher -and perhaps different distributions of- protein intake [31] were needed to enhance muscle strength and accretion of skeletal muscle mass. It should be also noted that the power calculation for sample size was based on lean mass as the primary outcome, rather than muscle strength. Therefore, it is unknown whether a larger sample size was needed to reveal a significant change in handgrip strength.

It has been previously reported that protein supplementation can enhance function in older adults. Namely, Tieland et al. [53] provided an oral supplement delivering 15 g of protein

twice daily (with breakfast and lunch) to older frail adults. This addition resulted in significantly enhanced physical performance. The potential for high protein meals to maintain or increase muscle mass and strength in older adults has been recently reported by Loenneke et al. [54] who showed that one or two meals containing 30-45 g protein · day⁻¹ were associated with higher lean mass and strength compared with those who did not consume any meals over the threshold of 30g protein.

4.2 Effects of exercise training on body composition and function in sarcopenic obesity

The mechanisms underpinning the effects of exercise on body composition and function in older age are mainly accounted for by regulation of genes, circulating hormone levels (e.g. testosterone, IGF-1) and metabolic pathways (especially by activating the mechanistic target of rapamycin (mTOR), which is a pathway also activated by leucine-rich protein meals [31]) and have been reviewed in detail by Garatachea et al. [55] and McGlory and Phillips [56]. However, it is still unclear whether the modulation of these pathways can translate into real-world benefits for adults with SO.

In study B[48], the aim was to assess the effect of high-speed resistance exercise training on indices of SO. In spite of possible methodological limitations, the improvements in strength, power and IADL reported in study B[42], provide some evidence that exercise can improve several domains of physical performance such as strength and power in older adults with SO. This is in agreement with previous reports supporting the benefits of resistance exercise training on clinically important outcomes, even in the absence of increased muscle mass [8][31][57]. This may be partly accounted for by the adaptive plasticity in the neuromuscular system and skeletal muscle tissue in response to resistance exercise even in advanced older

age [58]. A significant improvement particularly in power can be very important for individuals with SO since muscle power can be a predictor of mobility skills and a more influential indicator of physical capacity compared with absolute changes in strength [59]. Another interesting finding from study B[48] was the large effect size observed in peak leg power achieved by exercising at 50 % 1RM. To a certain extent, this finding may be explained by the novel aspect of the study design, that is, the resistance exercise progression protocol. Resistance load would increase only when a power plateau was reached [48]. Therefore, the protocol was designed in such a way as to favor maximum power output.

The lack of significant changes in lean mass or muscle mass after exercise training in adults with sarcopenia, (which has also been reported elsewhere [8]), may be accounted for by protocol-specific differences such as: duration, type, intensity, volume and frequency of exercise, as well as the availability of adequate nutrients (protein/amino acids), which are needed to elicit an anabolic response and consequently muscle hypertrophy [31]. One limitation of study B[48] was the lack of control for dietary intake, which could have partly explained the lack of effect on body composition. It has been shown that a bout of resistance exercise can stimulate muscle protein synthesis (MPS) to a higher degree than protein breakdown, however, in the absence of post-workout provision of nutrients (especially protein) it can result in negative net muscle protein balance [60][61], and is a limitation of study designs to date.

These data support the potential benefit of a resistance exercise program within lifestyle intervention protocols due to its positive effect on muscle strength, power and function in older adults with SO. Although no statistically significant body-composition changes were reported in the included studies, the significant improvements in strength, power and function

may be more important for the quality of life of adults with SO than absolute changes in body fat or lean mass per se.

4.3 Recommendations for future research

More intervention trials should be undertaken to identify effective lifestyle strategies in adults with SO, that will inform more robust approaches to combat this condition. Future research should also bridge the gap in knowledge with respect to multimodal approaches combining resistance exercise training with dietary strategies modulating protein intakes, in order to augment muscle mass and strength [32] or fat free mass [62], and potentially alongside an energy-deficit diet to promote fat loss [33].

It is important to note that although the need to augment muscle mass is paramount, a reduction in fat mass and especially fat infiltrating the muscle tissue is equally important, since intermuscular fat can result in mobility limitations [16]. Exercise training can preferentially reduce intermuscular adipose tissue more effectively than caloric-restriction alone [63]. However, a combination of exercise with caloric-restriction can lead to greater losses of total fat mass, which in turn may result in greater improvements in physical function, sometimes even at the expense of lean tissue [64][65]. Nevertheless, it is currently unknown whether this loss of lean mass may be detrimental in the long term for the life quality of an older individual with sarcopenia who has already experienced a large decline in muscle mass and strength.

To our knowledge only three studies to date, have reported significantly increased muscle or lean mass while concomitantly reducing fat mass [66-68]. The pilot study conducted by

Maltais et al. [68] was the only one that recruited older overweight adults with a low appendicular lean mass index. The authors concluded that 16-weeks of a whole body resistance exercise regimen (at 80% 1RM) followed by the consumption of ~13 g dairy supplement (chocolate milk with added milk powder) increased lean mass and reduced fat mass even at the absence of caloric restriction ($n=8$). In the same study, the group that received a soy-based beverage (matched for energy, protein, essential amino acids, carbohydrate and calcium content) ($n=8$), reported significant increases only in lean mass but no changes in fat mass. Similar results, but in pre-menopausal women, have been reported by Josse et al. [66]. After 16 weeks of mixed exercise training (aerobic and resistance) combined with a caloric restriction (500 kcal daily deficit), only the high protein group (30% of total energy intake came from protein, half of which was derived from dairy products) experienced lean gains concomitant with fat losses. Albeit in younger adults, a recent four-week intervention combining $2.4 \text{ g protein} \cdot \text{kg bw}^{-1} \cdot \text{day}^{-1}$ (achieved by consuming 3-4 whey-based protein beverages daily) with a mixed resistance, plyometric and high-intensity interval training alongside an energy deficit regimen resulted in significantly higher lean mass and lower body fat [67]. The control group which differed only in protein intake ($1.2 \text{ g protein} \cdot \text{kg bw}^{-1} \cdot \text{day}^{-1}$) did not experience a significant change in lean mass [67].

Although the effectiveness of the aforementioned paradigms needs to be evaluated in older adults with SO, they provide the framework for an initial approach to combat this condition. What is primarily lacking from the literature is trials recruiting older adults with SO. Additionally, protocols combining a high protein diet (potentially using whey or dairy proteins) with a modest caloric deficit along with a well structured exercise training regimen could be adopted. Moreover, long-term and follow-up studies with adults with SO who have

intentionally lost weight should be undertaken, in order to assess the impact of weight loss (especially if it comes from lean tissue) on life quality and other comorbidities.

4.4 Limitations

The main limitation of this review is the scarcity of data and studies undertaken in older groups with SO. We reviewed only studies with participants presenting the sarcopenic phenotype, as defined by the EWGSOP [1], using an appropriate methodology to assess body composition. Although the EWGSOP reached a consensus in 2010 on the definition and assessment of sarcopenia, adopting criteria for low muscle and low strength or function, there are recent studies [68][69] that use solely the criterion of low muscle mass to define sarcopenia as it was initially proposed [70], without taking into consideration low strength or function. In addition, one study conducted before 2010 was excluded because muscle mass was assessed using urinary creatinine [71], a method not included in the EWGSOP definition. Regarding the two studies included in this review, study B[48] aimed to recruit specifically participants with sarcopenic obesity. Study A[47] used sarcopenia as an inclusion criterion, and although the mean % BF values met our cut-off criteria for obesity, after personal communication with the authors it was reported that although the majority of participants had the sarcopenic obesity phenotype, a small proportion (5 out of 40) had a % BF below our cut-off.

Although of vital importance, there is an apparent lack of interventions with older adults with SO. This is in accordance with Finger et al. [62] who commented that interventions may refer to or discuss sarcopenia, however, the number of studies recruiting adults specifically with sarcopenia is very limited. This is even more complex with respect to studies on sarcopenic obesity, with a number of reviews [26][36][33][72] presenting interventions with participants

who either had none or only one of the conditions (sarcopenia or overweight/obesity but not both) and extrapolate these results to propose ways to improve the sarcopenic obesity phenotype. An example of an intervention study is that by Gadelha et al. [73] investigating the effect of exercise training on changes in the sarcopenic obesity index (assessed by the residuals method initially presented by Newman et al. [40] and adopted by the EWGSOP) of older Brazilian women. However, older age was the main inclusion criterion and no criteria specific for sarcopenia or obesity were adopted.

4.5 Conclusion

This review assessed studies investigating the effectiveness of exercise or nutritional interventions to improve the body composition and strength/function of older adults with sarcopenia and obesity. None of the included studies significantly reduced body fat or increased either skeletal muscle mass or lean mass. Although the number of included studies was low, it is evident that exercise training can elicit significant improvements in aspects of physical fitness such as muscle strength and power, and consequently improve performance in activities of daily living in adults with SO. The addition of 15 g protein·day⁻¹ to the habitual diet via cheese consumption revealed a non-statistically significant trend towards increased handgrip strength, and a significantly better insulin response in men, but not in women. The lack of published data highlights the necessity for new research adopting universally accepted cut-offs for sarcopenic obesity, with the inclusion of appropriately designed exercise programs and dietary regimens, and with detailed assessments of dietary patterns and protein intakes for the targeted population group.

Supplementary material

Supplementary file 1 (.pdf) **S1 Table**. Assessment of the methodological quality of the included studies with the modified Downs and Black Scale.

Supplementary file 2 (.pdf) **S1 Appendix**. Search strategy

Supplementary file 3 (.doc) **PRISMA Checklist**

Acknowledgements

We would like to thank Queen Margaret University's librarians and especially Sheila Williams for the guidance they provided during the initial stages of this systematic review.

Authors' contributions

CT, CAG, EB and JJ designed the study protocol and contributed to the writing of the manuscript. CT conducted the search and screening of titles, abstracts, full-text articles, the study selection, data extraction and quality assessment and prepared the manuscript. JJ screened the full-text articles and assessed the eligibility of the studies. CAG conducted the extraction, analysis and interpretation of data and risk of bias (quality) assessment. All authors read the final version of the manuscript.

Competing interests

The authors have no competing interests to declare

Funding

Queen Margaret University, Edinburgh provided support in the form of a studentship for the main author [CT], but did not have any role in the decision to produce and publish this manuscript or in the conceptualization, methodology, analysis and interpretation of data presented.

References

- [1] Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European working group on Sarcopenia in older people. *Age and Ageing* 2010;39:412–23. doi:10.1093/ageing/afq034.
- [2] Visser M, Schaap LA. Consequences of Sarcopenia. *Clinics in Geriatric Medicine* 2011;27:387–99. doi:10.1016/j.cger.2011.03.006.
- [3] Visser M, Newman AB, Nevitt MC, Kritchevsky SB, Stamm EB, Goodpaster BH, et al. Reexamining the Sarcopenia hypothesis: Muscle mass versus muscle strength. *Annals of the New York Academy of Sciences* 2000;904:456–61. doi:10.1111/j.1749-6632.2000.tb06500.x.
- [4] Newman AB, Kupelian V, Visser M, Simonsick EM, Goodpaster BH, Kritchevsky SB, et al. Strength, but not muscle mass, is associated with mortality in the health, aging and body composition study cohort. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences* 2006;61:72–7. doi:10.1093/gerona/61.1.72.
- [5] Wang X, Miller GD, Messier SP, Nicklas BJ. Knee strength maintained despite loss of Lean Body mass during weight loss in older obese adults with knee osteoarthritis. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences* 2007;62:866–71. doi:10.1093/gerona/62.8.866.
- [6] Clark BC, Manini TM. What is dynapenia? *Nutrition* 2012;28:495–503. doi:10.1016/j.nut.2011.12.002.
- [7] Gong Z, Muzumdar RH. Pancreatic function, type 2 diabetes, and metabolism in aging. *International Journal of Endocrinology* 2012;2012:1–13. doi:10.1155/2012/320482.

666 [8] Cruz-Jentoft AJ, Landi F, Schneider SM, Zuniga C, Arai H, Boirie Y, et al. Prevalence of
667 and interventions for sarcopenia in ageing adults: A systematic review. Report of the
668 international Sarcopenia initiative (EWGSOP and IWGS). *Age and Ageing* 2014;43:748–59.
669 doi:10.1093/ageing/afu115.

670 [9] WHO. Obesity and overweight. World Health Organization 2016.
671 <http://www.who.int/mediacentre/factsheets/fs311/en/> (accessed August 19, 2016).

672 [10] Porter Starr KN, McDonald SR, Bales CW. Obesity and physical Frailty in older adults:
673 A Scoping review of lifestyle intervention trials. *Journal of the American Medical Directors*
674 *Association* 2014;15:240–50. doi:10.1016/j.jamda.2013.11.008.

675 [11] Fakhouri TH, Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity among
676 older adults in the United States, 2007–2010. *NCHS Data Brief* 2012;106:1–8.

677 [12] Public Health England and Food Standards Agency. Official Statistics National Diet and
678 Nutrition Survey: Results from Years 1 to 4 (combined) of the Rolling Programme for 2008
679 and 2009 to 2011 and 2012 NDNS Chapter 4 Tables, Table 41a 2014.
680 https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/311001/NDNS
681 [_Appendices_and_tables.zip](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/311001/NDNS) (accessed June 17, 2016).

682 [13] Houston DK, Ding J, Nicklas BJ, Harris TB, Lee JS, Nevitt MC, et al. Overweight and
683 obesity over the adult life course and incident mobility limitation in older adults: The health,
684 aging and body composition study. *American Journal of Epidemiology* 2009;169:927–36.
685 doi:10.1093/aje/kwp007.

686 [14] Himes CL, Reynolds SL. Effect of obesity on falls, injury, and disability. *Journal of the*
687 *American Geriatrics Society* 2011;60:124–9. doi:10.1111/j.1532-5415.2011.03767.x.

688 [15] Oreopoulos A, Kalantar-Zadeh K, Sharma AM, Fonarow GC. The obesity paradox in the
689 elderly: Potential mechanisms and clinical implications. *Clinics in Geriatric Medicine*
690 2009;25:643–59. doi:10.1016/j.cger.2009.07.005.

691 [16] Visser M, Goodpaster BH, Kritchevsky SB, Newman AB, Nevitt M, Rubin SM, et al.
692 Muscle mass, muscle strength, and muscle fat infiltration as predictors of incident mobility
693 limitations in well-functioning older persons. *The Journals of Gerontology Series A:*
694 *Biological Sciences and Medical Sciences* 2005;60:324–33. doi:10.1093/gerona/60.3.324.

695 [17] Greenberg AS, Obin MS MS. Obesity and the role of adipose tissue in inflammation and
696 metabolism. *Am J Clin Nutr* 2006;83:461S – 465S.

697 [18] Visser M, Pahor M, Taaffe DR, Goodpaster BH, Simonsick EM, Newman AB, et al.
698 Relationship of Interleukin-6 and tumor necrosis factor- with muscle mass and muscle
699 strength in elderly men and women: The health ABC study. *The Journals of Gerontology*
700 *Series A: Biological Sciences and Medical Sciences* 2002;57:M326–32.
701 doi:10.1093/gerona/57.5.m326.

702 [19] Koster A, Ding J, Stenholm S, Caserotti P, Houston DK, Nicklas BJ, et al. Does the
703 amount of fat mass predict age-related loss of Lean Mass, muscle strength, and muscle
704 quality in older adults? *The Journals of Gerontology Series A: Biological Sciences and*
705 *Medical Sciences* 2011;66A:888–95. doi:10.1093/gerona/qlr070.

706 [20] Waters DL, Baumgartner RN. Sarcopenia and obesity. *Clinics in Geriatric Medicine*
707 2011;27:401–21. doi:10.1016/j.cger.2011.03.007.

708 [21] Baumgartner RN. Body composition in healthy aging. *Annals of the New York*
709 *Academy of Sciences* 2006;904:437–48. doi:10.1111/j.1749-6632.2000.tb06498.x.

710 [22] Baumgartner RN, Wayne SJ, Waters DL, Janssen I, Gallagher D, Morley JE. Sarcopenic
711 obesity predicts instrumental activities of daily living disability in the elderly. *Obesity*
712 *Research* 2004;12:1995–2004. doi:10.1038/oby.2004.250.

713 [23] Rolland Y, Lauwers-Cances V, Cristini C, van Kan GA, Janssen I, Morley JE, et al.
714 Difficulties with physical function associated with obesity, sarcopenia, and sarcopenic-
715 obesity in community-dwelling elderly women: The EPIDOS (EPIDemiologie de
716 l'OSteoporose) study. *American Journal of Clinical Nutrition* 2009;89:1895–900.
717 doi:10.3945/ajcn.2008.26950.

718 [24] Tian S, Xu Y. Association of sarcopenic obesity with the risk of all-cause mortality: A
719 meta-analysis of prospective cohort studies. *Geriatrics & Gerontology International*
720 2015;16:155–66. doi:10.1111/ggi.12579.

721 [25] Prado CMM, Wells JCK, Smith SR, Stephan BCM, Siervo M. Sarcopenic obesity: A
722 critical appraisal of the current evidence. *Clinical Nutrition* 2012;31:583–601.
723 doi:10.1016/j.clnu.2012.06.010.

724 [26] Poggiogalle E, Migliaccio S, Lenzi A, Donini LM. Treatment of body composition
725 changes in obese and overweight older adults: Insight into the phenotype of sarcopenic
726 obesity. *Endocrine* 2014;47:699–716. doi:10.1007/s12020-014-0315-x.

727 [27] Batsis JA, Mackenzie TA, Barre LK, Lopez-Jimenez F, Bartels SJ. Sarcopenia,
728 sarcopenic obesity and mortality in older adults: Results from the national health and
729 nutrition examination survey III. *European Journal of Clinical Nutrition* 2014;68:1001–7.
730 doi:10.1038/ejcn.2014.117.

731 [28] Naseeb MA, Volpe SL. Protein and exercise in the prevention of sarcopenia and aging.
732 *Nutr Res* 2017;04:1-20. doi: <http://dx.doi.org/10.1016/j.nutres.2017.01.001>

733 [29] Phillips SM. Nutritional supplements in support of resistance exercise to counter age-
734 related Sarcopenia. *Advances in Nutrition: An International Review Journal* 2015;6:452–60.
735 doi:10.3945/an.115.008367.

736 [30] McLeod M, Breen L, Hamilton DL, Philp A. Live strong and prosper: The importance of
737 skeletal muscle strength for healthy ageing. *Biogerontology* 2016;17:497–510.
738 doi:10.1007/s10522-015-9631-7.

739 [31] Witard OC, McGlory C, Hamilton DL, Phillips SM. Growing older with health and
740 vitality: A nexus of physical activity, exercise and nutrition. *Biogerontology* 2016.
741 doi:10.1007/s10522-016-9637-9.

742 [32] Cermak NM, Res PT, de Groot LC, Saris WH, van Loon LJ. Protein supplementation
743 augments the adaptive response of skeletal muscle to resistance-type exercise training: A
744 meta-analysis. *American Journal of Clinical Nutrition* 2012;96:1454–64.
745 doi:10.3945/ajcn.112.037556.

746 [33] Weinheimer EM, Sands LP, Campbell WW. A systematic review of the separate and
 747 combined effects of energy restriction and exercise on fat-free mass in middle-aged and older
 748 adults: Implications for sarcopenic obesity. *Nutrition Reviews* 2010;68:375–88.
 749 doi:10.1111/j.1753-4887.2010.00298.x.

750 [34] Bauer J, Biolo G, Cederholm T, Cesari M, Cruz-Jentoft AJ, Morley JE, et al. Evidence-
 751 based recommendations for optimal dietary protein intake in older people: a position paper
 752 from the PROT-AGE Study Group. *J Am Med Dir Assoc.* 2013;14:542-59.

753 [35] Deutz NE, Bauer JM, Barazzoni R, Biolo G, Boirie Y, Bosy-Westphal A, et al. Protein
 754 intake and exercise for optimal muscle function with aging: recommendations from the
 755 ESPEN Expert Group. *Clin Nutr.* 2014;33:929-36.

756 [36] Goisser S, Porzel S, Volkert D, Kemmler W, Sieber C, Bollheimer C, et al. Sarcopenic
 757 obesity and complex interventions with nutrition and exercise in community-dwelling older
 758 persons – a narrative review. *Clinical Interventions in Aging* 2015:1267.
 759 doi:10.2147/cia.s82454.

760 [37] Phillips SM. Determining the protein needs of "older" persons one meal at a time. *Am J*
 761 *Clin Nutr.* 2017;105:291-2.

762 [38] Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic
 763 reviews and meta-analyses: The PRISMA statement. *International Journal of Surgery*
 764 2010;8:336–41. doi:10.1016/j.ijssu.2010.02.007.

765 [39] Nelson ME, Rejeski WJ, Blair SN, Duncan PW, Judge JO, King AC, et al. Physical
 766 activity and public health in older adults: recommendation from the American College of
 767 Sports Medicine and the American Heart Association. *Medicine & Science in Sports &*
 768 *Exercise* 2007;39:1435–45. doi:10.1249/mss.0b013e3180616aa2.

769 [40] Newman AB, Kupelian V, Visser M, Simonsick E, Goodpaster B, Nevitt M, et al.
 770 Sarcopenia: Alternative definitions and associations with lower extremity function. *Journal of*
 771 *the American Geriatrics Society* 2003;51:1602–9. doi:10.1046/j.1532-5415.2003.51534.x.

772 [41] Downs SH, Black N. The feasibility of creating a checklist for the assessment of the
 773 methodological quality both of randomised and non-randomised studies of health care
 774 interventions. *Journal of Epidemiology & Community Health* 1998;52:377–84.
 775 doi:10.1136/jech.52.6.377.

776 [42] Eng J, Teasell R, Miller W, Wolfe D, Townson A, Aubut J-A, et al. Spinal cord injury
 777 rehabilitation evidence: Method of the SCIRE systematic review. *Topics in Spinal Cord*
 778 *Injury Rehabilitation* 2007;13:1–10. doi:10.1310/sci1301-1.

779 [43] Olivo SA, Macedo LG, Gadotti IC, Fuentes J, Stanton T, Magee DJ. Scales to assess the
 780 quality of Randomized controlled trials: A systematic review. *Physical Therapy*
 781 2007;88:156–75. doi:10.2522/ptj.20070147.

782 [44] Saunders LD, Soomro GM, Buckingham J, Jamtvedt G, Raina P. Assessing the
 783 methodological quality of Nonrandomized intervention studies. *Western Journal of Nursing*
 784 *Research* 2003;25:223–37. doi:10.1177/0193945902250039.

785 [45] Wouters-Wesseling W. The effect of a liquid nutrition supplement on body composition
786 and physical functioning in elderly people. *Clinical Nutrition* 2003;22:371–7.
787 doi:10.1016/s0261-5614(03)00034-7.

788 [46] Kwon J, Yoshida Y, Yoshida H, Kim H, Suzuki T, Lee Y. Effects of a combined
789 physical training and nutrition intervention on physical performance and health-related
790 quality of life in Prefrail older women living in the community: A Randomized controlled
791 trial. *Journal of the American Medical Directors Association* 2015;16:263.e1–263.e8.
792 doi:10.1016/j.jamda.2014.12.005.

793 [47] Aleman-Mateo H, Macias L, Esparza-Romero J, Astiazaran-Garcia H, Blancas AL.
794 Physiological effects beyond the significant gain in muscle mass in sarcopenic elderly men:
795 Evidence from a randomized clinical trial using a protein-rich food. *Clinical Interventions in*
796 *Aging* 2012;225. doi:10.2147/cia.s32356.

797 [48] Balachandran A, Krawczyk SN, Potiaumpai M, Signorile JF. High-speed circuit training
798 vs hypertrophy training to improve physical function in sarcopenic obese adults: A
799 randomized controlled trial. *Experimental Gerontology* 2014;60:64–71.
800 doi:10.1016/j.exger.2014.09.016.

801 [49] Pennings B, Groen B, de Lange A, Gijsen AP, Zorenc AH, Senden JMG, et al. Amino
802 acid absorption and subsequent muscle protein accretion following graded intakes of whey
803 protein in elderly men. *AJP: Endocrinology and Metabolism* 2012;302:E992–9.
804 doi:10.1152/ajpendo.00517.2011.

805 [50] Yang Y, Breen L, Burd NA, Hector AJ, Churchward-Venne TA, Josse AR, et al.
806 Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey
807 protein in older men. *British Journal of Nutrition* 2012;108:1780–8.
808 doi:10.1017/s0007114511007422.

809 [51] Moore DR, Churchward-Venne TA, Witard O, Breen L, Burd NA, Tipton KD, et al.
810 Protein ingestion to stimulate Myofibrillar protein synthesis requires greater relative protein
811 Intakes in healthy older versus younger men. *The Journals of Gerontology Series A:*
812 *Biological Sciences and Medical Sciences* 2014;70:57–62. doi:10.1093/gerona/glu103.

813 [52] Tieland M, Borgonjen-Van den Berg KJ, van Loon LJC, de Groot LCPGM. Dietary
814 protein intake in community-dwelling, frail, and institutionalized elderly people: Scope for
815 improvement. *European Journal of Nutrition* 2011;51:173–9. doi:10.1007/s00394-011-0203-
816 6.

817 [53] Tieland M, van de Rest O, Dirks ML, van der Zwaluw N, Mensink M, van Loon LJ, et
818 al. Protein supplementation improves physical performance in frail elderly people: a
819 randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc.* 2012;13:720-6.

820 [54] Loenneke JP, Loprinzi PD, Murphy CH, Phillips SM. Per meal dose and frequency of
821 protein consumption is associated with lean mass and muscle performance. *Clinical Nutrition*
822 2016;35:1506–11. doi:10.1016/j.clnu.2016.04.002.

823 [55] Garatachea N, Pareja-Galeano H, Sanchis-Gomar F, Santos-Lozano A, Fiuza-Luces C,
824 Moran M, et al. Exercise attenuates the major hallmarks of aging. *Rejuvenation Res.*
825 2015;18:57-89.

826 [56] McGlory C, Phillips SM. Exercise and the Regulation of Skeletal Muscle Hypertrophy.
827 Prog Mol Biol Transl Sci. 2015;135:153-73.

828 [57] Churchward-Venne TA, Tieland M, Verdijk LB, Leenders M, Dirks ML, de Groot
829 LCPGM, et al. There are no Nonresponders to resistance-type exercise training in older men
830 and women. Journal of the American Medical Directors Association 2015;16:400–11.
831 doi:10.1016/j.jamda.2015.01.071.

832 [58] Aagaard P, Suetta C, Caserotti P, Magnusson SP, Kjaer M. Role of the nervous system
833 in sarcopenia and muscle atrophy with aging: Strength training as a countermeasure.
834 Scandinavian Journal of Medicine & Science in Sports 2010;20:49–64. doi:10.1111/j.1600-
835 0838.2009.01084.x.

836 [59] Bean JF, Leveille SG, Kiely DK, Bandinelli S, Guralnik JM, Ferrucci L. A comparison
837 of leg power and leg strength within the InCHIANTI study: Which influences mobility more?
838 The Journals of Gerontology Series A: Biological Sciences and Medical Sciences
839 2003;58:M728–33. doi:10.1093/gerona/58.8.m728.

840 [60] Phillips S, Tipton K, Aarsland A, Wolf S, Wolfe R. Mixed muscle protein synthesis and
841 breakdown after resistance exercise in humans. The American Journal of Physiology
842 1997;273:E99–107.

843 [61] Koopman R, van Loon LJC. Aging, exercise, and muscle protein metabolism. Journal of
844 Applied Physiology 2009;106:2040–8. doi:10.1152/jappphysiol.91551.2008.

845 [62] Finger D, Goltz FR, Umpierre D, Meyer E, Rosa LHT, Schneider CD. Effects of protein
846 Supplementation in older adults undergoing resistance training: A systematic review and
847 Meta-Analysis. Sports Medicine 2014;45:245–55. doi:10.1007/s40279-014-0269-4.

848 [63] Murphy JC, McDaniel JL, Mora K, Villareal DT, Fontana L, Weiss EP. Preferential
849 reductions in intermuscular and visceral adipose tissue with exercise-induced weight loss
850 compared with calorie restriction. J Appl Physiol (1985). 2012;112:79-85.

851 [64] Villareal DT, Chode S, Parimi N, Sinacore DR, Hilton T, Armamento-Villareal R, et al.
852 Weight loss, exercise, or both and physical function in obese older adults. N Engl J Med.
853 2011;364:1218-29.

854 [65] Beavers KM, Beavers DP, Nesbit BA, Ambrosius WT, Marsh AP, Nicklas BJ, et al.
855 Effect of an 18-month physical activity and weight loss intervention on body composition in
856 overweight and obese older adults. Obesity (Silver Spring). 2014;22:325-31.

857 [66] Josse AR, Atkinson SA, Tarnopolsky MA, Phillips SM. Increased consumption of dairy
858 foods and protein during diet- and exercise-induced weight loss promotes fat mass loss and
859 lean mass gain in overweight and obese premenopausal women. J Nutr. 2011;141:1626-34.

860 [67] Longland TM, Oikawa SY, Mitchell CJ, Devries MC, Phillips SM. Higher compared
861 with lower dietary protein during an energy deficit combined with intense exercise promotes
862 greater lean mass gain and fat mass loss: A randomized trial. American Journal of Clinical
863 Nutrition 2016. doi:10.3945/ajcn.115.119339.

864 [68] Maltais ML, Perreault K, Courchesne-Loyer A, Lagacé JC, Barsalani R, Dionne IJ.
865 Effect of resistance training and various sources of protein Supplementation on body fat mass

866 and metabolic profile in Sarcopenic overweight older adult men: A pilot study. *International*
867 *Journal of Sport Nutrition and Exercise Metabolism* 2016;26:71–7. doi:10.1123/ijsnem.2015-
868 0160.


869 [69] Mason C, Xiao L, Imayama I, Duggan CR, Foster-Schubert KE, Kong A, et al. Influence
870 of diet, exercise, and serum vitamin D on Sarcopenia in Postmenopausal women. *Medicine &*
871 *Science in Sports & Exercise* 2013;45:607–14. doi:10.1249/mss.0b013e31827aa3fa.

872 [70] Evans WJ. What is Sarcopenia? *The Journals of Gerontology Series A: Biological*
873 *Sciences and Medical Sciences* 1995;50A:5–8. doi:10.1093/gerona/50a.special_issue.5.

874 [71] Yarasheski K, Pak-Loduca J, Hasten D, Obert K, Brown M, Sinacore. Resistance
875 exercise training increases mixed muscle protein synthesis rate in frail women and men ≥ 76
876 yr old. *The American Journal of Physiology* 1999;277:E118–25.


877 [72] Li Z, Heber D. Sarcopenic obesity in the elderly and strategies for weight management.
878 *Nutrition Reviews* 2012;70:57–64. doi:10.1111/j.1753-4887.2011.00453.x.

879 [73] Gadelha AB, Paiva FML, Gauche R, de Oliveira RJ, Lima RM. Effects of resistance
880 training on sarcopenic obesity index in older women: A randomized controlled trial. *Archives*
881 *of Gerontology and Geriatrics* 2016;65:168–73. doi:10.1016/j.archger.2016.03.017.



882

883 **Figure 1.** Relationship between sarcopenia and obesity and associated risks as well as
 884 management strategies. Notes: Solid arrow: direct and positive association; Dashed line
 885 management strategy attenuating/reversing the condition:



886

887 **Figure 2. Information flow through the phases of the systematic review according to**
 888 **PRISMA guidelines.**

889 **Table 2. Summary key points of the included study designs.**

Study	Summary	Strengths	Limitations
Study A Aleman-Mateo et al. [47]	<ul style="list-style-type: none"> ○ Nutritional Intervention ○ 40 participants ○ 3 months ○ Habitual diet plus 210g Ricotta cheese ·day⁻¹ (intervention) vs habitual diet (control) 	<ul style="list-style-type: none"> ○ Intention to treat analysis ○ Body composition by dual-energy xray absorptiometry ○ Physically-independent participants. ○ Baseline and follow up clinical tests for kidney and liver function ○ Blinded personnel delivering the assessment tests. 	<ul style="list-style-type: none"> ○ Lack of baseline and follow up dietary intake and physical activity data
Study B Balachandran et al. [48]	<ul style="list-style-type: none"> ○ Exercise Intervention ○ 21 participants ○ 15 weeks ○ High speed circuit resistance (HSC) training (intervention) vs strength hypertrophy (SH) resistance training (control) 	<ul style="list-style-type: none"> ○ Independent living community-dwellers. ○ Participants were partially blinded to the intervention. ○ Testing personnel blinded ○ All sessions supervised by 2 physiology majors 	<ul style="list-style-type: none"> ○ No allocation concealment ○ Per-protocol analysis ○ Underpowered ○ Characteristics of participants lost to follow-up not described ○ No description of the exercise setting

890 **Table 2. Summary of the included studies**

Study	Setting/ Study Design/ Duration	Group	Participants Mean Age (SD)/ characteristics	Exercise Training	Nutritional Intervention	Sample Size (n) Drop-out (DO n) Female (F n) Adherence (%)	Assessment of a)body composition b) strength or function	Outcome Measure
Study A (Aleman-Mateo et al. [47])	Mexico/ RCT: two arms, one control, one intervention / 3 months	Control	76.7 (5.8) / physically-independent, sarcopenic based on low TASM and strength, obese based on %BF	No	Habitual diet (HD)	Baseline n=20, Final n= 12 DO n= 3 F n=12 N/A	a) DXA b) HG strength	TASM→, FM→, LM→, HG→
		Intervention	75.4 (5.0)/ independent living sarcopenic based on low TASM and strength, obese based on %BF	No	HD plus 210 g of ricotta cheese/day, (providing 15.7 gr extra protein/day)	Baseline n=20, Final n=17 DO n=8 F n=11 N/A		TASM→, FM→ , LM→, HG→
Study B (Balachandran et al. [48])	USA/ RCT: Two arms, one control, one intervention/ 15 weeks	Control	71 (8.2)/ independent living community dwellers from South Miami, sarcopenic based on SMI and strength, obese based on %BF and BMI	Strength-hypertrophy (SH) training, 11 exercises, 3 sets of 10-12 reps per set at 70% 1RM	No	Baseline n=10, Final n=9 DO n= 1 F n=8 85%	a) BIA b) HG strength, SPPB, Leg press 1RM, Chest press 1RM, Leg press power, Chest press power,	SMI→, %BF→, SPPB→, Leg 1RM↑**, Leg Power↑*, Chest 1RM↑*, Chest Power↑**, HG→
		Intervention	71.6 (7.8)/ independent living community dwellers from South Miami	High speed circuit (HSC) training, 11 exercises: 3 circuits of 10-12 reps per exercise at loads that maximised peak power output	No	Baseline n=11, Final n= 8 DO n= 3 F n =8 81%		SMI→, %BF→, SPPB↓*, Leg 1RM→, Leg Power↑**‡, Chest 1RM↑**, Chest power↑**, HG→

891 Notes: → no significant change, ↑significant increase, ↓significant decrease, * <0.05 , ** <0.01 , ‡ significantly better than the control group;
892 %BF, percent body fat; BMI, body mass index; DXA, dual-energy xray absorptiometry; FM, fat mass; HG, handgrip; LM, lean mass; RCT, randomised control Trial; RM,
893 repetition maximum; SPPB, short physical performance battery test; TASM, total appendicular skeletal muscle.

